

# Comparisons of AIRS profiles against dedicated sondes (work in progress)


Bill Irion, Eric Fetzer,  
Van Dang, Evan Manning

Jet Propulsion Laboratory  
California Institute of Technology


With many thanks to all who provided  
sonde data to us

# Sonde data characteristics

- ~880 sondes
- ~7400 AIRS matchups within 3 hrs and 100 km



# Modus operandi



# AIRS –Sonde comparisons made

- L2 result compared to sonde profile interpolated to AIRS support grid
  - By level if temperature
  - By slab column if gas

- Init profile to sonde
  - How good (or bad) was our first guess

- L2 to “Kerned” sonde profile:  $\hat{x}$ 
  - Sonde profile multiplied by averaging kernel:

$$\hat{x} = x_{init} + A(x_{sonde} - x_{init})$$

- This is what AIRS “should have seen,” given its sensitivity and vertical resolution

# Sample results

All qual\_temp flags  $\leq 1$   
2 hr, 100 km matchup range


Only one AIRS observation per  
sonde added to average

# of  
matchups

"Unkerned"

"Kerned"


Initial



# Heredia, Costa Rica


## Temperature profile comparison by season

Only one AIRS observation per sonde added to average  
All temperature qual flags  $\leq 1$ , within 100 km, 2 hrs



# Comparison for Chesapeake SON season by cloud fraction

One-to-one matchup, all temperature qual flags  $\leq 1$ , within 100 km, 2 hrs



# Further work

- Testing & debugging
- Quality control for sondes
  - E.g., handling data dropouts
- Water vapor (in progress), ozone
- Validation against different seasons, cloud conditions etc.
- Validation against different climate regimes
- Results to be used in V5 validation report

# **PRELIMINARY INTERCOMPARISON OF UT/LS TEMPERATURE PROFILES BETWEEN AIRS, GPS, AND ECMWF (abridged)**

**Baijun Tian<sup>1,2</sup>**

**V. Dang<sup>2</sup>, F. Irion<sup>2</sup>, E. Fetzer<sup>2</sup>, J. Teixeira<sup>2</sup>, C. Ao<sup>2</sup>, B. Wilson<sup>2</sup>, G. Mani<sup>2</sup>**

<sup>1</sup>Joint Institute for Regional Earth System Science and Engineering (JIFRESSE), UCLA

<sup>2</sup>Jet Propulsion Laboratory (JPL), California Institute of Technology (Caltech)

**AIRS Science Team Meeting, Oct 2008, Greenbelt, MD**

# UT/LS


1. **The upper troposphere and lower stratosphere (UT/LS) is the important layer responsible for the troposphere-stratosphere exchange.**
2. **Accurate knowledge of tropopause temperature, pressure and height is very important for detecting the global climate change.**

## OBJECTIVE

**To examine the ability of AIRS temperature retrieval to delineate the tropopause.**



# MATCHUP LOCATIONS FOR JAN 2003

< 100 km, < 2 hr





Black Squares - GPS; Red Asterisks - AIRS

# WESTERN PACIFIC



# ICELAND



# **SUMMARY**

- 1. AIRS can roughly capture the tropopause with an error of maybe 20hPa although detailed comparisons need to be done.**
- 2. There are significant differences between AIRS and GPS temperature profiles (2-4K) especially near the tropopause.**
- 3. There is a strong correlation between AIRS and ECMWF temperature profile errors relative to GPS.**